



All Databases



Enter Text Here

Recent searches



Go to PDB


Top page

Protein


Ligands

Prot-pr

Periplasmic binding protein



Main view



Bottom view



Right view



Contents

Description

[Header details](#)[Header records](#)[References](#)[PROCHECK](#)

Protein chains

229 a.a.

Ligands

FLC x2

Waters x349

Tools

[Image Generation](#)[AstexViewer™@MSD-EBI](#)[Run PROCHECK](#)[Clefts Calculation](#)

PDB id: 2hxw

Name: Periplasmic binding protein

Title: Crystal structure of peb3 from campylobacter jejuni

Structure: Major antigenic peptide peb3. Chain: a, b. Engineered: ye

Source: Campylobacter jejuni. Strain: nctc 11168. Gene: peb3. Ex
escherichia coli.

UniProt: Chains A, B: Q0PBL7 (Q0PBL7_CAMJE) [Pfam]

Seq: 250

Struc: 229

Key: PfamA domain Secondary structure

Resolution: 1.60Å

R-factor: 0.188

R-free: 0.212

Authors: E.S.Rangarajan,S.Bhatia,D.C.Watson,C.Munger,M.Cygler
A.Matte,N.M.Young,Montreal-Kingston Bacterial Structural
Initiative (Bsgi)Key ref: E.S.Rangarajan et al. (2007). Structural context for protein
glycosylation in bacteria: The structure of PEB3, an adhesin
Campylobacter jejuni.. *Protein Sci*, 16, 990-995.

[PubMed id: 17456748] [DOI: 10.1110/ps.062737507]

Date: 04-Aug-06

Release date: 01-May-07

DOI no: [10.1110/ps.062737507](https://doi.org/10.1110/ps.062737507)
 PubMed id: [17456748](https://pubmed.ncbi.nlm.nih.gov/17456748/)

Key
Protein Sci 16:99

Structural context for protein N-glycosylation in bacteria: The structure of PEB3, an adhesin from *Campylobacter jejuni*.

E.S.Rangarajan, S.Bhatia, D.C.Watson, C.Munger, M.Oygier, A.Matte, N.M.Young.

ABSTRACT



Campylobacter jejuni is unusual among bacteria in possessing a eukaryotic-like system for N-linked protein glycosylation residues in sequons of the type Asp/Glu-Xaa-Asn-Xaa-Ser/Thr. However, little is known about the structural context of glycosylated sequons, limiting the design of novel recombinant glycoproteins. To obtain more information on sequon structure we have determined the crystal structure of the PEB3 (Cj0289c) dimer. PEB3 has the class II periplasmic-binding protein with each monomer having two domains with a ligand-binding site containing citrate located between them, and overall resembles molybdate- and sulfate-binding proteins. The sequon around Asn90 is located within a surface-exposed loop of two structural elements. The three key residues are well exposed on the surface; hence, they may be accessible to the oligosaccharyltransferase in the folded state.

Selected figure(s)




Figure 2.

Figure 2. PEB3-binding site and sequon structure. (A) Citrate binding-site density, 2F0-FC (omit) electron density map contour.

The above figure is reprinted by permission from the Protein Society: Protein Sci (2007, 16, 990-995) copyright 2007. Figure was selected by an automated process.

Literature references that cite this PDB file's key reference

PubMed id Reference

- [18316380](https://pubmed.ncbi.nlm.nih.gov/18316380/) T.D.Ho, B.M.Davis, J.M.Ritchie, and M.K.Walder (2008).
 Type 2 secretion promotes enterohemorrhagic *Escherichia coli* adherence and intestinal colonization.
Infect Immun, 76, 1858-1865.

The most recent references are shown first. Citation data come partly from CiteXplore and partly from an automated harvest. Note that this is likely to be only a partial list as not all journals are covered by either method. However, we are continuously updating citation data so more and more references will be included with time.

[Terms of Use](#) [EBI Funding](#) [Contact EBI](#) © European Bioinformatics Institute 2006-2008. EBI is an Outstation of the European Molecular Biology Laboratory
